

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE
1995

3. REPORT TYPE AND DATES COVERED

Proceedings of DoD Conference Oct. 24-26, 1995

4. TITLE AND SUBTITLE

Retrieval of Cirrus Radiative and Spatial Properties Using Coincident Multispectral Imager and Sounder Satellite Data

5. FUNDING NUMBERS

N/A

6. AUTHOR(S)

R.P.d'Entremont, D.P.Wylie, Szu-Cheng Ou and Kuo-Nan Liou

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Atmospheric and Environmental
Research, Inc.
840 Memorial Dr.
Cambridge, MA 02139-3794

Department of Meteorology/CARSS
University of Utah
Salt Lake City, UT 84112

Space Science and Engineering Center
University of Wisconsin
Madison, WI 53706

8. PERFORMING ORGANIZATION
REPORT NUMBER

N/A

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

SERDP
901 North Stuart St. Suite 303
Arlington, VA 22203

10. SPONSORING / MONITORING
AGENCY REPORT NUMBER

N/A

11. SUPPLEMENTARY NOTES

Proceedings of the Cloud Impacts on DoD Operations and Systems 1995 Conference (CIDOS-95). No copyright is asserted in the United States under Title 17, U.S. code. The U.S. Government has a royalty-free license to exercise all rights under the copyright claimed herein for Government purposes. All other rights are reserved by the copyright owner.

12a. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release: distribution is unlimited

12b. DISTRIBUTION CODE

A

13. ABSTRACT (Maximum 200 Words)

In addition to the wide variability in properties common for the other types of clouds, cirrus clouds have transmissivity values t_λ that span the entire possible domain $0 \leq t_\lambda \leq 1$. This variability adds complexity to the analysis of cirrus clouds. In comparison to opaque clouds, uncertainties exist in thin cirrus cloud amount, altitude, thickness, and optical properties as retrieved from satellite because the measured cirrus signal is affected additionally by an unknown radiation component from below.

Cirrus radiative and spatial properties are derived using HIRS CO₂ Slicing and multispectral AVHRR imager techniques. Each of these models is based on radiative transfer principles that intrinsically account for both the semi-transparent nature of thin cirrus clouds and the attenuation effect of atmospheric water vapor in the MWIR and LWIR thermal window regions.

14. SUBJECT TERMS

SERDP, t_λ , transmissivity, AVHRR, HIRS CO₂ Slicing, attenuation

15. NUMBER OF PAGES

2

16. PRICE CODE

N/A

17. SECURITY CLASSIFICATION
OF REPORT

unclass

18. SECURITY CLASSIFICATION
OF THIS PAGE

unclass

19. SECURITY CLASSIFICATION
OF ABSTRACT

unclass

20. LIMITATION OF ABSTRACT

UL

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

CLOUD IMPACTS ON DOD OPERATIONS AND SYSTEMS 1995 CONFERENCE

U.S. Air Force Phillips Laboratory - Science Center
Hanscom Air Force Base, Massachusetts
24-26 October 1995



19980806 069

CIDOS - 95

Cloud Modeling and Data for Defense Simulation Activities
"Emphasizing Sufficient Physical Reality in Simulating Clouds"

PL-TR-95-2129
Environmental Research Papers, No. 1179

**PREPRINT OF THE CLOUD IMPACTS ON
DoD OPERATIONS AND SYSTEMS
1995 CONFERENCE (CIDOS-95)**

Editor

Donald D. Grantham

1 October 1995

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED



PHILLIPS LABORATORY
Directorate of Geophysics
Air Force Materiel Command
Hanscom Air Force Base, MA 01731-3010

Retrieval of Cirrus Radiative and Spatial Properties Using Coincident Multispectral Imager and Sounder Satellite Data

Robert P. d'Entremont

Atmospheric and Environmental Research, Inc.
840 Memorial Drive, Cambridge, MA 02139-3794

Donald P. Wylie

Space Science and Engineering Center
University of Wisconsin, Madison, Wisconsin 53706

Szu-Cheng Ou and Kuo-Nan Liou

Department of Meteorology / CARSS
University of Utah, Salt Lake City UT 84112

In addition to the wide variability in properties common for other types of clouds, cirrus clouds have transmissivity values t_λ that span the entire possible domain $0 \leq t_\lambda \leq 1$. This variability adds complexity to the analysis of cirrus clouds. In comparison to opaque clouds, uncertainties exist in thin cirrus cloud amount, altitude, thickness, and optical properties as retrieved from satellite because the measured cirrus signal is affected additionally by an unknown radiation component from below.

Cirrus radiative and spatial properties are derived using HIRS CO₂ Slicing and multispectral AVHRR imager techniques. Each of these models is based on radiative transfer principles that intrinsically account for both the semi-transparent nature of thin cirrus clouds and the attenuation effect of atmospheric water vapor in the MWIR and LWIR thermal window regions.

Comparison is made of cirrus cloud attributes, both spatial and radiative, obtained for the same cloud scene using measurements from the independent AVHRR and HIRS sensors onboard the NOAA polar orbiting satellites. While the fundamental requirement is the same for both the HIRS and AVHRR models, i.e. to detect the presence of thin cirrus and to determine its radiative and spatial attributes, the capabilities of the two techniques depart from each other in

numerous respects. The most important distinctions are based on the differences in the spectral bandpass and spatial resolutions of the HIRS and AVHRR sensors.

Three cirrus retrieval algorithms, CO₂ Slicing, AVHRR, and SERCAA have been shown to complement one another for increasing the accuracy of the obtained cirrus parameters. New techniques will be presented that combine the strongest and most reliable attributes of the imager and sounder-based cirrus retrieval algorithms in combination with SERCAA-derived background analyses to generate an improved overall cirrus analysis. The "background" may be either the clear ground or an underlying water-droplet cloud. Improved SERCAA estimates of underlying cloud and surface temperature can significantly improve the AVHRR and HIRS CO₂ Slicing determination of cirrus bulk emissivity.

The SERCAA-derived cirrus fraction N is useful to CO₂ slicing in separating the effect of N from the effective emissivity $N\epsilon$ for those cirrus clouds whose optical attributes are uniform within a particular HIRS FOV, thus allowing for direct comparisons of the true thermal infrared cirrus bulk emissivities retrieved by each algorithm. On the other hand, the CO₂ slicing technique provides an independent determination of cirrus effective altitude z which obviates the need for the AVHRR parameterization of the variation of ϵ with wavelength.

Better cirrus emissivity analyses in turn will significantly improve the accuracy of cirrus radiative models.